

Presents:

DRUG LAB UNLIMITED

Advanced Rules for Narcotics Use and Creation

Written by Gary Ostlelford for Cyberpunk 2020, (ocelot@cox.net), Edited for Interlock Unlimited by Deric Bernier (droc@mc2k.com), Additional materiel written by Seth Skorkowsky



This is just one of many articles for the Interlock Unlimited System you can find on Datafortress 2020, the largest, most comprehensive Cyberpunk 2020 site in the world.

Visit it at:

http://datafortress2020.oliwy.net/

UPDATED 02-10-12



DRUG LAB 101

DRUG CREATION

Let's Jump Right Into It, Shall We?

When creating a drug, you must ask some questions:

- 1.) What does the drug do? What benefits does the drug grant the user? Why would anyone want to take such a drug?
- 2.) What sorts of negative side effects are there? Do these effects occur after taking the drug, or when it wears off?
- 3.) How strong is the drug? Is it powerful or mild? How easy is it to overdose on this drug?
- 4.) How long do the effects last?
- 5.) Is the drug illegal, or can you buy it at any drug or liquor store?
- 6.) What form does the drug come in? Do you swallow it, snort it, or shoot it?

With these optional rules, you can create new and interesting chemicals to cripple your characters with. The basic formula from the standard rules still applies. You choose the Drug Effect(s) you want, and add the costs together. Next, you choose any Side Effects that want your drug to have. Side effects reduce the base difficulty number. After side effects, you choose the drug's Strength, which adds from between 1 and 16 to the base difficulty, and then you choose its duration, which multiplies the base difficulty by 1, 2, or 3 times (the longer the drug lasts, the harder it is to produce).

Once all the numbers have been assimilated, you have the drug's base difficulty. A base difficulty can never be a negative amount, and at the very least must equal 1. If, by chance, you end up with a negative number, drop the negative and use it as if it were positive (drugs that really mess you up aren't all that easy to make, either). If the base difficulty is less than 10, the Pharmacist must still roll a minimum of a 10 on their skill check to create it, however they still go by the Actual Difficulty for pricing to sell and create.

The base difficulty multiplied by the drug's legality modifier will give you the drug's base cost. The last step you take is to choose what form the drug comes in. The drug's form will affect its final street cost.

In the Beginning, man created drugs. And for a while, they were good. That is, until they wore off...

DRUG EFFECTS -

People take drugs for any number of reasons. One of the most common among these is to "escape." The drug makes you feel good, and for a while you forget your problems. It's a form of relaxation, like a chemical massage. But there's also more to it than that. Some people take drugs to give them that extra something, that special "edge" that makes them just that much better. While much of this can be psychological, some of it can be attributed to a drug's effect on the person's physiology.

Below are listed certain drug effects, what happens if you OD (see OVERDOSE) while taking one, and what they mean to the discerning pharmaceutical gourmet. Base difficulties are listed in parentheses. Note that a drug effect can only be bought ONCE.

Antibiotic (Difficulty 10)

These drugs prevent infection, or stop an existing one. While I could fill an entire file on microbiology, and the effects of antibiotics on germs, this would do very little for game play. Assume that an antibiotic, when employed against an infection that is susceptible to it, will cause the infection to cease spreading and the user to become well more quickly, where the strength of the drug is the multiplier. For example, someone with strep throat who takes a +2 STR antibiotic will get well twice as fast, and so on.

OD: Overdosing on antibiotics will cause severe nausea and diarrhea for 1d6 days.

Anticoagulant (Difficulty 10)

The opposite of coagulants, Anticoagulants prevent blood from clotting. While usable as a quick antidote from a coagulant overdose, such drugs can also be employed to allow an affected user to bleed to death. Assume that someone who has used anticoagulants and is subsequently wounded for four or more points of damage will lose an additional point per turn until he dies from blood loss or the bleeding is stopped.

OD: Extreme hemophilia will result in the case of anticoagulant overdose. Even the smallest cuts (1 point of damage or more) will bleed heavily, and the user will suffer 1 point of damage per turn until he exsanguinates or the wound is stanched.

Antidote (Difficulty 15)

Antidotes add their strength to saves made against poisons, toxins, etc. Normally, these drugs are tailored to reduce the effects of one specific substance, or negate them altogether.

OD: Most antidotes are almost poisons in their own right, and anyone overdosing on them will suffer the effects of minor poisoning. Treat this poisoning as the drug's STR in D6 poison damage. For example, overdosing on a +3 STR antidote would cause 3D6 damage to the user.

Aphrodisiacs (Difficulty 10)

These drugs increase a person's sex drive and libido. The user will be physically uncomfortable unless involved in a sexual situation, and in the case of some stronger drugs, this may not even help unless the user is continually stimulated. While under the influence, the user has a negative modifier to seduction checks equal to the drug strength.

OD: The character's sexual urges cannot be satisfied, and his state of arousal will not diminish for up to several days or weeks.

Attribute Increase (Difficulty 20)

This increases an attribute by the drug's strength rating for the drug's duration. Almost any attribute can be increased. Such attributes include Reflex, Intelligence, Body, Movement Allowance, Empathy, Cool, and Tech.

OD: If an overdose occurs, roll 1d10. On a roll of 1-5, you have lost a point from the attribute in question, permanently.

Coagulant (Difficulty 10)

These drugs cause blood to clot, and prevent blood loss. In games where blood loss is a genuine concern, assume that a patient's blood loss is reduced by as many points as the drug has in STR. In other cases where blood loss does not apply, the drug STR may be

added to a First Aid or Medtech roll when stabilizing a patient.

OD: An overdose of coagulants can cause blood to clot inside a healthy body, unavoidably causing death. Make a death save as if you were at Mortal 2, with a -4 modifier. Failure indicates death.

Contraceptive (Difficulty 10)

These drugs prevent pregnancy. They have a base effectiveness of 80%, plus 5% for every point of drug strength.

OD: An overdose of contraceptive drugs can cause sterility. Roll 1d10. On a roll of 1, the user is sterile and cannot sire children.

Depressant (Difficulty 5)

Depressants slow you down. They reduce tension, stress, and help you relax. The down side is that they also reduce your Reflex score by an amount equal to the drug's strength. While you may wonder why someone might want to take a drug like this, remember that alcohol is a depressant, and it's the most popular drug around.

OD: Taking too many depressants will cause the character to make a stun/shock at -2 or slip into a deep sleep/coma for 1d10 hours.

Enhanced Perception (Difficulty 15)

The user of these sorts of drugs starts to notice the most intricate of details. While this adds the drug's strength to his Awareness score for the duration, the user may be overcome by his increased sensory powers. Loud noises, bright lights, intense tastes, and strong odors have twice the normal effect on him.

OD: Anyone suffering the effects of overdosing on Enhanced Perception drugs will be completely overwhelmed by his senses, and cannot perform any actions. There is a 25% chance that he will suffer from terrible hallucinations.

Euphoric (Difficulty 5)

These drugs make you feel really, REALLY good. They give a "rush" that lasts for as long as the drug's duration.

OD: Overdosing on a euphoric generally makes you very sick, including nausea, cold sweats, and either sleeplessness or sleepiness.

Hallucinogen (Difficulty 10)

Wow, man, look at the butterflies. These drugs make the user see things that aren't there. While under the influence, or "tripping," the user is prone to see anything his frying imagination can cook up. Such hallucinations depend on the mood of the subject (as well as the GM!), but aren't usually terrible unless the drug itself is bad.

OD: If overdosed on, assume that the hallucinogenic "trip" goes completely bad. In the case of extremely traumatic hallucinations, mental illness may result (ie, phobias, etc.).

Hypnotic (Difficulty 15)

Hypnotic drugs, in high doses, are used to produce sleep during surgery. In low doses, they make many users speak and talk less guardedly about things they might otherwise have trouble speaking about. Due to this side-effect, hypnotics are routinely used as "truth serums" by corporate, military, and government agencies. In game terms, hypnotic drugs reduce the user's COOL by their strength, while at the same time granting a bonus equal to the drug's strength to anyone using Interrogation, Human Perception, or Interview on the subject.

OD: Hypnotics, if taken in large doses, cause a deep anesthesia for 1d10 hours.

Increased Endurance (Difficulty 10)

These drugs include most stimulants. While under the influence of these types of drugs, the user has increased energy and stamina. This is reflected by adding the drug's strength to his Endurance skill checks. However, someone under the influence of such a drug may have trouble sleeping.

OD: In the case of an OD, the user must make a stun/shock save at -2. In the case of a failure, he suffers from a heart attack, and must then make a death save as if at Mortal 0 or die.

Increased Healing Rate (Difficulty 15)

These drugs speed up the body's natural healing process by .5 points per point of drug strength per day. Available only by prescription.

OD: If healing drugs are taken to the point of overdose, assume that the user enters a severe state of shock as the body attempts to heal nonexistent wounds and injuries, and must make a death save at Mortal 0 or die.

Pain Negation (Difficulty 10)

Pain killers, plain and simple. These drugs allow the user to ignore wound penalties by one level per strength point the drug has. For example, someone who is seriously wounded and takes a +1 STR pain killer acts as if he is only lightly wounded, and so forth.

OD: The user must make a successful unmodified stun/shock save or slip into a coma for 1d10 hours.

Soporific (Difficulty 5)

Soporifics put you to sleep. Otherwise known as sleeping pills or tranquilizers. A user who wishes to resist the effects of these drugs must roll a body check using the strength of the drug as a negative modifier. Failure indicates immediate unconsciousness for the duration of the drug. Success means that the user gets a negative modifier equal to the drug's STR to ALL actions until the drug wears off.

OD: We all know what happens when you take too many sleeping pills. You go to sleep and never wake up. Assume that without proper treatment, someone who has ODed on soporifics will die (no death save is given) or enter a deep coma which will last for 1d10 days.

Stun Reducers (Difficulty 10)

Like pain killers, but these are usually taken before one is actually wounded. They allow the user to gain a bonus to his stun/shock and death saves, +1 for each point of drug strength. People under the influence of powerful stun-reducing drugs have been known to walk through a shower of high caliber bullets as if pebbles were being rained upon them. Of course, most of them died later on.

OD: In the case of most stun reducing drugs, an overdose will cause the user to become extremely hostile and enter a psychotic rage akin to the side effect (see SIDE EFFECTS).

Toxin: Biologic (Difficulty 10)

Bio-Toxins are poisons that attack the body's organs, causing them to cease function and possible death. Bio-Toxins cause 2D6 points of damage per level to the user. A successful BOD check using the strength of the drug as a negative modifier reduces the damage to 1D6 points of damage for every level.

Toxin: Neurologic (Difficulty 15)

Neuro-Toxins are poisons that attack the body's nervous system. Synapses in the brain and body either cease communication with organs, or go into overdrive, causing lethal seizures. In both cases, Nerve Toxins cause 2D10 points of damage per level to the user. A successful BOD check using the strength of the drug as a negative modifier reduces the damage to 1D10 points of damage for every level

DRUG FEATURES

Drug Features are beneficial modifications that can be worked into drugs. They add a significant amount of difficulty, but the benefits they grant are sometimes worth it. Difficulty Cost Modifiers are listed in parentheses. Note that a drug feature can only be taken once per drug.

Lessened Withdrawal Symptoms (Cost Modifier +10)

Only usable for addictive drugs. Withdrawal symptoms are half as bad as they would normally be. A drug with the "Severe Withdrawal Symptoms" side effect cannot contain this feature.

Longer Duration (Cost Modifier +10)

When the drug is taken, roll the duration and then double the result.

Timed Duration (Cost Modifier +10)

Drug duration lasts as long as is specified in the description, but no longer than the maximum duration. For instance, a drug with a medium duration can last anywhere from ten to sixty minutes, but no longer. Drugs with this side effect usually require exact dosages based on the weight of the subject. Note that giving a subject too much of a drug with this feature will produce an immediate overdose, while giving a subject too little will have no effect at all.

AND NOW FOR SOMETHING COMPLETELY DIFFERENT - SIDE EFFECTS!

Side effects are the bad things that drugs do to you. They come in two varieties - UNTIMED and TIMED.

Untimed side effects are the big, bad ones. They happen as soon as the drug is imbibed, and cause permanent damage to the user of the drug. Each Untimed effect can only be used in a drug's creation once

There are two types of Timed side effects: IMMEDIATE and DELAYED. Immediate side effects manifest as soon as the drug takes effect, while Delayed side effects occur after the drug wears off (ie, "hangover"). When choosing a Timed side effect for a drug, you must specify if the effect you are adding to your cocktail is either Immediate or Delayed. Note that the same side effect can be taken twice, one for an Immediate effect and one for a Delayed one. Immediate effects last until the drug wears off. Delayed effects kick in as soon as the drug wears off, and last as long as the drug's previously rolled duration.

Timed side effects can be bought as "possible side effects" at 1/2 cost, dropping fractions. A possible side effect only occurs on a 1D10 roll of 1-5. For instance, if "Kidney/Liver Failure" is bought as a possible side effect, it will grant only a -4 cost modifier (instead of -8), and its effects will only be felt 50% of the time. The list of side effects follows. Difficulty Cost Modifiers are listed in parentheses. Note that stat reductions are cumulative.

UNTIMED SIDE EFFECTS

Addiction Tolerance (Cost Modifier -4)

People who are addicted to the drug gain a tolerance to its effects. While still hooked on the drug, they gain no benefit from it unless they double or triple their dosage. Even when the dosage is upped, users only gain benefits equal to those normally indicated for a second (or third) dose (which is to say, half), but they are still at risk of overdose.

Carcinogenic (Cost Modifier -10)

The drug causes cancer in laboratory tests, and the Surgeon General recommends that you stick a loaded gun in your mouth as opposed to taking the substance. Every time you use the substance, roll 1D100 (or have your GM do it for you, secretly). On a roll of 01-03, you've got a little bitty tumor growing somewhere in your body. Of course, tumors continue to spread and grow, don't they...

Cardiac Arrest/Heart Attack (Cost Modifier -12)

This particularly deadly side effect indicates that the drug puts massive strain on the user's heart, risking a severe heart attack. Roll 1D10. On a result of 1-3, cardiac arrest results. Roll another 1D10, and then make a death save as if you were at the resulting Mortal level. For example, if the second 1D10 roll results in a 6, the user makes a death save as if at Mortal 6. Failure indicates death.

Coma (Cost Modifier -10)

By using this drug, characters risk falling into a deep coma. When the drug is taken, roll 1D10. On a result of 1-3, the character becomes comatose for 1D10 days per strength point of the drug. Unless properly cared for, death will most likely result from starvation and dehydration.

Death (Cost Modifier -15)

The drug that this side effect is attached to is no better than poison. Each time you use it, you must make a Death Save as if you were at Mortal 2. Failure indicates your heart stops and you die. Pretty simple, eh?

Genetic Damage (Cost Modifier -6)

Drugs with this side effect screw with a person's genes. In practical terms, this can result in severe birth defects among the user's progeny. Make a note every time the drug is taken. If the user ever manages to sire a child, roll 1D10. If the roll is less than or equal to the number of times the drug has been taken by the user in his lifetime, there may be some sort of defect/mutation in his child. To determine the defect, use the "Offspring Mutation Table" on page 22 of "Deep Space," treating all rolls of "Favorable" as "Harmless."

Nerve Degeneration (Cost Modifier -15)

This side effect is reserved for those really nasty drugs that only idiots and guinea pigs imbibe of. Every time you use this drug, your Reflex stat is reduced by 2. If it ever reaches 1, you've contracted a serious case of Parkinson's Disease, and all you can do is drool. At least you can use those neat handicapped parking spaces now. If it goes to 0 or below, roll up a new character, and remember to say no to drugs.

Nightmares (Cost Modifier -6)

This drug causes terrible nightmares in users. The first time a user of such a drug sleeps after a dose, he must roll under his COOL -4. Failure indicates that he wakes up screaming after 1D6 hours of restless slumber, and will be unable to sleep again for hours

afterwards. Nightmares caused by the side effect cannot be recalled by the user, and reflect on his most primal fears. Addicts of such drugs often use them in conjunction with stimulants since they know that monsters are waiting for them behind their closed eyelids...

Permanent Sense Loss (Cost Modifier -6)

The drug causes the user to lose a sense (touch, smell, hearing, taste, or sight) permanently. Roll 1D10. On a roll of 1-3, the affected sense is gone, forever. This is due to nerve damage, so senses enhanced with cybernetics (ie, optics, hearing modules, tactile boosts, etc.) are affected as well. It is possible to regain the sense with expensive nanosurgery involving nerve reconstruction.

Physiologically Addictive (Cost Modifier -5/-10/-15)

These drugs are physiologically addictive. A drug with this side effect causes the user to physically require the drug in order to function. Like "Psychologically Addictive", this side effect has three costs which reflect on how addictive the chemical in question is. -5 is Slightly Addictive, -10 is Mildly Addictive, and -15 reflects that the drug is Highly Addictive.

Possible Attribute Loss (Cost Modifier -8)

This side-effect covers the possible loss of an attribute from a variety of factors which could result from drug ingestion. Attributes that can be damaged this way include Intelligence and Tech (brain damage), Reflex and Movement Allowance (nerve degeneration), Body (immunodeficiency and muscle atrophy), and Empathy and Cool (psychological damage). Every time you take the drug in question, roll 1d10. On a roll of 1-3, you've lost a point from one of these attributes permanently.

Psychologically Addictive (Cost Modifier -4/-8/-12)

This side effect reflects the psychological effect the drug has on its user. A user who is psychologically addicted to a substance is convinced he needs the drug, and feels he can't perform without it. This side effect has three costs, reflecting on how ddictive the substance in question is. -4 is Slightly Addictive, -8 is Mildly Addictive, and -12 reflects that the drug is Highly Addictive.

Severe Withdrawal Symptoms (Cost Modifier -6)

Only usable for addictive drugs. Withdrawal symptoms are twice as bad as they would normally be. A drug with the "Lessened Withdrawal Symptoms" feature cannot contain this side effect.

Sterility (Cost Modifier -8)

Using this drug can cause the junkie to become unable to sire children by damaging his sexual glands directly. In severe cases, sexual function is completely impaired, and impotence will result. Roll 1d10 each time the drug is used. On a roll of 1-3, the user's sexual fertility is destroyed.

TIMED SIDE EFFECTS -

Acne (Cost Modifier -2)

The drug causes the user's face to break out due to excess oil production or stress. Normally, this could start out as a small problem, but frequent use of the drug can cause a more serious acne problem to develop (negative modifiers to ATT, as well as the increased chance of infection).

Aggressive Behavior (Cost Modifier -6)

A character experiencing such behavior will act to satisfy his own needs with little regard for the feelings of others, is irritable and easily provoked. He will not back down from a fight, and may

even attempt to initiate one. He will have no respect for authority, and any suggestion that he is incapable of anything will be seen as a challenge.

Bad Breath (Cost Modifier -2)

This side effect indicates that halitosis develops. The character's breath will be incredibly rank, granting him a -3 modifier to all social rolls involving direct, face-to-face contact. No amount of brushing or Listerine will help.

Bloodshot Eyes (Cost Modifier -2)

The drug induces red, puffy, bloodshot eyes. Vision is not impaired.

Cold Sweats (Cost Modifier -2)

This side effect causes the character to sweat for no apparent reason. No amount of anti-perspirant will do, since every pore on his body is working a double shift. Sweaty characters suffer no real modifications, though they may grow dehydrated or get the chills in cold or air-conditioned environments (-1 to all checks involving Reflex). Their palms will be slippery, and they will stink. Incredibly sweaty characters may also incur negative modifiers to social rolls.

Constipation (Cost Modifier -2)

Constipation prevents a character from performing normal bowel movements for the duration of the side effect. No matter how bad the character feels he has to go, no amount of straining, grunting, or groaning will help. The character will feel as if he is carrying a lead weight in his colon, and will receive a -1 modifier to all actions involving physical exertion, including Initiative rolls.

Cowardice (Cost Modifier -6)

The character will exhibit his most timid side, and will be more likely to run from trouble. Any time the character tries to accomplish a task which may cause him physical harm, he must make a Save against Cool at -2 in order to show some guts and stick around. If he fails, he must run or hide, whichever is more convenient.

Dandruff/Eczema (Cost Modifier -2)

The user's skin cracks and peels away. This side effect could be confined to the user's scalp, a specific part of the body, or the body as a whole. Complications include reduced Attractiveness. In addition, the user tends to leave little bits of himself behind at crime scenes, and police will have no problem isolating his DNA.

Dehydration (Cost Modifier -2)

The subject cannot retain water at all, and will suffer from symptoms such as cotton mouth and dry eyes. This side effect is usually accompanied by either nausea, diarrhea, diuretic, cold sweats, or a combination of the four.

Delusions (Cost Modifier -5)

Anyone suffering from delusions is under the impression that something that is highly unlikely is actually happening. Such delusions can be as minor as believing that Abraham Lincoln was controlled by aliens, or as serious as believing that you actually ARE Abraham Lincoln.

Depression (Cost Modifier -4)

The character becomes depressed, and doesn't feel like doing anything. Anything he attempts to do will suffer a -2 modifier, because he just doesn't really care. Severe cases of depression can cause feelings of hopelessness, and characters who are normally

unstable may feel like dropping the final curtain by committing suicide.

Diarrhea (Cost Modifier -4)

The drug causes sudden, painful, and explosive diarrhea. The unlucky soul will be unable to control his bowels for more than a few moments at a time.

Difficult Respiration (Cost Modifier -6)

Anyone experiencing the effect of difficult respiration is suffering from either shortness of breath, asthma-like symptoms, a tightening of the diaphragm muscles, or a combination of the three. On a roll of 1 on 1D10 indicates the subject cannot, in fact, breathe at all, and will suffocate unless proper medical attention has been administered. Running is out of the question, as is most other high-and low-impact exercise.

Diuretic (Cost Modifier -2)

The drug increases the amount of urine produced and secreted by the body. The user affected by this will exhibit excessive urination, at least once per hour of the effect's duration, if not more.

Dizziness (Cost Modifier -4)

The drug causes severe dizziness. The character suffering this effect will have trouble walking in a straight line or, for that matter, standing. This results in the user's Reflex and MA being halved for the duration of the side effect. In addition, he looks damn weird as he lurches around, bumping into objects and people.

Easily Bruised (Cost Modifier -2)

Even small bumps and thumps leave large bruises on the user. These bruises tend to be painful and noticeable, but don't affect the user otherwise.

Excessive Salivation (Cost Modifier -2)

The drug causes the user's salivary glands to work double time. Unless he continues to swallow his saliva as it is produced, it will dribble down his chin and onto his body in an incredibly disgusting fashion. This side effect can also be used to emulate the "mad dog" effect of foaming at the mouth.

Flatulence (Cost Modifier -2)

The user gets terrible abdominal gas, and must vent uncontrollably for the duration. While being mildly painful, the real disadvantage to this side effect comes with the stench it causes, as well as the potential embarrassment. Then again, there are people who enjoy this sort of thing, so who am I to judge?

Hair Loss (Cost Modifier -2)

The drug causes hair loss in both men and women. Eighty or ninety percent of the user's head hair falls out effortlessly in large clumps. Most users of the drug are left with a mangy-looking scalp and small patches of unhealthy-looking hair here and there, modifying the character's ATT by -2 or more.

Hallucinations (Cost Modifier -5)

This side effect causes unlooked-for hallucinations. Such hallucinations can be traumatic or harmless, depending on the situation. Any Hallucinogenic drug with this side effect causes a "bad trip" 95% of the time.

Headaches (Cost Modifier -4)

The drug causes blinding headaches. The character cannot focus on anything for long periods of time, and has trouble seeing. Loud noises and bright lights will make this side effect worse.

Heartburn/Acid Indigestion (Cost Modifier -2)

The drug causes an excess production of stomach acids. While this is uncomfortable and rarely serious, it can result in ulcers in the stomach and esophagus.

Hives (Cost Modifier -4)

The character develops itchy and painful hives all over his body. Not only does he suffer from the "Itchy" side-effect, but the have-related rash covering his body is obvious and unattractive, as well as being incredibly uncomfortable.

Impotence/Frigidity (Cost Modifier -2)

While the user might desire to have sex, he or she is physically incapable for the duration of the side effect.

Increased Pain Sensitivity (Cost Modifier -6)

The drug makes little pains feel much, much bigger. Big pains will incapacitate the character as he writhes around in agony. While this effect lasts, the character makes all stun/shock saves at -2, and gains a -4 to resist torture.

Insomnia (Cost Modifier -4)

The character cannot sleep, no matter how tired he is. Characters suffering from lack of sleep tend to be cranky and unable to concentrate. Subtract -1 from all actions until the character gets at least six hours of sleep.

Internal Bleeding (Cost Modifier -4)

For some reason, the drug causes the user's guts to hemorrhage and bleed. For every two hours of side effect duration (or every fraction thereof), the user takes one box of damage. Blood may show up in the users stools, urine, and/or vomit.

Irrational Fear (Cost Modifier -10)

Anyone suffering this side effect will develop a sudden intense fear of a random object, thing, or circumstance, determined by the Game Master. The character will stop at nothing to put as much distance between himself and the object/situation he fears. If escape is impossible, he will fight like a cornered animal.

Itchy (Cost Modifier -3)

The character suffers from a terrible itchy sensation all over his body. No amount of scratching or hydrocortisone will stop the irritation. Due to trouble concentrating, the character is at -1 to all activities. Habitual users of drugs that cause itching are usually walking scabs.

Kidney/Liver Failure (Cost Modifier -8)

Taking this drug will cause a user's kidneys and liver to work overtime while filtering his blood stream of the nasty substance. This can cause these organs to fail. Roll 1d10. On a roll of 1, the user's kidneys and/or liver have stopped functioning. Unless proper medical attention is sought, the character will die within twenty-four hours due to poisons running unchecked through his bloodstream.

Lack Of Concentration (Cost Modifier -5)

The character cannot concentrate on any one thing for more than a couple minutes. His mind is restlessly wandering, and he is easily distracted. For the duration, he suffers a -3 to all actions.

Light Sensitivity (Cost Modifier -2)

The drug dilates the pupils and induces red, bloodshot eyes. The character is at a loss in any situation involving a bright light (-2 to all actions in daylight) unless wearing sunglasses or equipped with

flare compensation. Characters with cybereyes are still affected by bright lights, since this side effect focuses on the optic nerve rather than the eye itself.

Loss Of Appetite (Cost Modifier -2)

The character will not eat, and will not feel hungry. People hooked on drugs with this side effect tend to be thin and emaciated. Characters who don't eat will starve within a couple weeks if not given treatment. This is reflected in the loss of one BODY point every three or four days. When the character's BODY reaches 0, he will die. If normal food consumption is resumed, the character will recover one BODY point a day until he reaches his original score.

Loss Of Inhibition (Cost Modifier -4)

Anyone suffering from Loss of Inhibition will do things that they normally would be adverse to doing. For instance, someone who wouldn't normally consider raping a helpless teenage girl might think it's a good idea at the time and throw inhibitions to the wind. Much of this side effect must be role-played. If game mechanics enter into it, however, the player must roll under his character's COOL score on a 1D10 in order to consider his actions and stop. Modify the character's COOL score by the drug's STR, using the number as a negative (instead of positive) modifier.

Memory Loss (Cost Modifier -4)

The user will have no memory of what has happened while under the effects of the drug. If he tries really hard (Very Difficult Intelligence check), he can remember vague events, but no specifics (ie, names, phone numbers, conversations, etc).

Nausea (Cost Modifier -4)

The character must make a stun/shock save at -2 or throw up. The nausea is painful and will extend into dry heaves. For the duration the character may not safely eat or drink anything. If he attempts to eat anything, a new stun/shock save at -2 is in order. Failure indicates more vomiting.

Numbness (Cost Modifier -4/-6)

At a -4 cost modifier, this drug causes the user's extremities to become numb and void of feeling. Fine manipulation is impossible, and walking is impossible unless the character is given support. At -6, the lack of feeling extends to the whole body. Since the user is experiencing no pain, the GM should secretly keep track of his wounds.

Paralysis (Cost Modifier -6)

Characters suffering from paralysis cannot move, speak, and they may have trouble breathing. Paralysis can take many forms, from complete muscle relaxation to a painful tightening of the entire muscular system. While paralyzed, the user is completely helpless and can perform no actions.

Paranoia (Cost Modifier -5)

While this side effect lasts, the user is convinced that everyone and everything is out to get him. While most of this will have to be role-played, a few Cool rolls at appropriate times may be called for.

Psychotic Rage (Cost Modifier -10)

A result of some of the worst combat drugs. The user goes berserk, attacking the nearest friend or enemy, he doesn't really care which. If a Difficult Resist Torture/Drugs roll is made, the user may specifically ignore a friend or ally and focus his attention on a target of his choice.

Reduced Attribute (Cost Modifier -5)

This drug reduces an attribute by an amount equal to the drug strength for an amount of time equal to the duration. Attributes that can be reduced include Reflex, Intelligence, Body, Movement Allowance, Empathy, Cool, and Tech. If a physical attribute (Reflex, Body, MA) is reduced below zero, make a Death Save or die. If a mental attribute (Intelligence, Tech) is reduced below zero, the user slips into a deep coma for the duration. In the case of Empathy being lowered below zero, assume that user is sociopathic and is liable to do anything. A Cool score below zero indicates total nervous breakdown.

Reduced Awareness (Cost Modifier -6)

The user of a drug with this effect is oblivious to the world around him. While Sense Reduction only affects one of the senses, Reduced Awareness affects them ALL. All skill rolls which rely on any of the user's senses suffer a -4 modifier, and all Awareness rolls are made at -6.

Ringing In Ears (Cost Modifier -2)

Due to severe tinnitus, the user has trouble hearing soft to medium volume sounds. As such, he suffers -2 to all noise-related awareness rolls.

Runny Nose (Cost Modifier -2)

Users suffering from a runny nose are constantly sniffling and blowing their noses. This doesn't really help, though. Roll 1d10. On a roll of 1 or 2, the character develops a bloody nose. While having no real game effect, he will be dripping blood everywhere. It will take between three and five minutes of constant pressure to stop the nose from bleeding, but the character still has a serious case of the sniffles.

Sense Reduction (Cost Modifier -4)

The drug reduces the user's sense of touch, smell, hearing, taste, or sight for the duration of the side effect. All rolls made which rely on the sense in question receive a -4 modifier. For example, someone suffering from Sense Reduction: Sight would be at a -4 to shooting his weapon due to blurred vision or near-blindness. All awareness rolls made with the appropriate sense are also at -4.

Sexual Arousal (Cost Modifier -4)

The drug turns the user on, yet to a lesser degree than most aphrodisiacs. Anyone attempting to seduce the user gains a +1 bonus. In addition, all outward signs of arousal are visible and quite obvious. If the drug with this side effect is an aphrodisiac, roll 1d10. On a result of 1 or 2, a permanent state of arousal occurs, and doesn't go away...EVER. Is that a truncheon in your pocket, or are you just happy to see me?

Sleepy (Cost Modifier -4)

The drug makes the user sleepy. A character who is overcome with sleepiness must make a successful stun/shock save or fall into a deep, dreamless sleep. Soporifics with this side effect induce a deep, coma-like state that can last for 1d10 days.

Strange Skin Pallor (Cost Modifier -2)

The drug tints the user's skin a strange color. This can indicate that the drug makes its user's skin pale, or that his lips turn blue, but it can also indicate a more severe chromatic variation to his epidermis. Like, say, your whole body is blue. Like a smurf.

Suicidal Tendencies (Cost Modifier -5)

The user becomes depressed or agitated, convinced that life is meaningless and that it might as well end as soon as possible. While possessed of suicidal tendencies, a character must roll under his COOL score -2. If the roll is failed, he must take immediate suicidal action. The method he uses is whichever is most convenient, be it putting a gun in his mouth and pulling the trigger, or jumping off his fifth floor balcony into the moving traffic below. Needless to say, users of such drugs eventually do society a favor.

Technicolor Excreta (Cost Modifier -2)

The user's bodily waste, be it urine or feces or both, turns a strange color. This effect lasts twice as long as the drug duration.

The Munchies (Cost Modifier -2)

The drug with this side effect causes the user to grow really irrationally hungry. The character will eat any edible substance, and will believe that it's the best thing he's ever tasted.

Tremors (Cost Modifier -2)

The user experiences violent tremors in his face and hands. The palsied fellow has his Reflex reduced by 2.

Water Retention (Cost Modifier -2)

Due to nonfunctional kidneys or other less traumatic problems, the user of a drug with this side effect will retain water for the side effect's duration. He will not be able to urinate, and will become bloated for the duration.

Weight Gain (Cost Modifier -2)

The drug causes the user's body to store excess energy in the form of fat. Users of drugs with this side effect tend to be quite large and overweight, while addicts suffer from obesity. Every time a drug with this effect is used, the user gains 1D6 pounds. Due to the extra weight he is carrying, an overweight character suffers a -1 to MA for every twenty-five pounds he weighs past his "ideal weight." In addition to the physical effects, people might suffer a social stigma.

DRUG STRENGTH

Not all drugs are created equal. Some are less potent than others, while some are dangerously strong. Hence, you have Drug Strength. The strength of a drug determines how powerful its effects are, how much can be taken before an overdose occurs, and how easy (or hard) they are to get hooked on if they are addictive. Drug strength can be any number between 1 and 5. The higher the number, the stronger the drug and the greater its effects.

Add the number chosen for the drug's strength rating to the base difficulty.

DRUG STRENTH	COST
1	1
2	2
3	4
4	8
5	16

ADDICTION

Some drugs are addictive. Someone who is addicted to a drug is physically or mentally dependent on it. Without their drug, they cannot operate at full capacity. When a drug is used, an addiction save may be called for. This depends entirely on how addictive the drug is (see the entries for PSYCHOLOGICAL and PHYSIOLOGICAL ADDICTION side effects).

How often an addiction save is called for depends on how addictive the drug is. Slightly Addictive drugs require an addiction

save every 1D6+4 times they are used, Mildly Addictive drugs every 1D3+2 times they are used, and Highly Addictive drugs require a roll each and every time they are used.

I recommend that the GM roll this number and make a note of it secretly. That way, players will only have a general idea how many times they can take a drug safely before they're forced to roll for addiction. If such a roll is called for, the user must roll above the drug's Addiction Number on 1D10. The addiction number of a drug is equal to it's strength times 1.75 rounding fractions to the nearest whole number. Addiction numbers are listed below:

DRUG STRENGTH	ADDICTION NUMBER
1	2
2	4
3	5
4	7
5	9

CRAVINGS AND WITHDRAWAL

Once a drug has worn off (see DRUG DURATION, below), an addict will need to get another dose. However, he won't necessarily need one right away. The chart below details the time a user can go before he is incapacitated by his need for the drug. When the user's most recent dose has worn off, roll the requisite number of dice under TIME TO NEXT DOSE to determine how long it will be before his cravings for the drug return.

If at that time he doesn't get a fix, roll the number of dice under TIME TO WITHDRAWAL to determine when, exactly, withdrawal symptoms will appear. It is recommended that the GM roll these values and keep them secret.

	TIME TO NEXT DOSE	TIME TO WITHDRAWAL
+1	1D6 Hours	5D6 Hours
+2	2D6 Hours	4D6 Hours
+3	3D6 Hours	3D6 Hours
+4	4D6 Hours	2D6 Hours
+5	5D6 Hours	1D6 Hours

Anyone attempting to kick a psychological addiction must stop taking the drug altogether. Withdrawal symptoms include -2 to all actions until withdrawal is complete (about a week or so). A strong psychological craving for the drug will remain, and a character who was once hooked may need to make a Cool roll (Resist Torture/Drugs applies) to resist the temptation if offered the drug again.

Physiological addiction is much stronger and harder to break. Kicking such an addiction takes about two weeks of treatment. All of the character's physical and mental attributes are reduced by half to reflect on the severe mental and physical anguish of the process. As with psychological addiction, the character will still crave the drug, and the same Cool roll applies if he comes into temptation's way.

TAKING MULTIPLE DOSES

Users may take multiple doses of a drug, but the drug's effects will not be doubled. In fact, for each subsequent dose after the first, halve all beneficial effects the drug grants, rounding fractions down. All negative side effects are not halved, however, making this a risky proposition. For example, a drug that causes death as a

side effect taken twice will cause two subsequent death saves to be made. While it may seem safe to take less powerful (and dangerous) drugs like they were M&M's, there is still the topic of Overdose.

OVERDOSE (O.D.)

When someone takes multiple doses of a drug, he increases his risk of becoming sick by introducing too much of the chemical into his body. If the total strength of combined drug doses is ever greater than 10 plus the character's BTM score (assume the BTM is a positive number), the character has a chance to become terribly ill from overdosing. The effects of an overdose vary greatly, depending on the drugs taken. Check the individual Drug Effects entries to find out exactly what happens to the user of the drug. If a drug has more than one effect (ie, a drug that acts as both a Euphoric and a Stun Reducer), apply both sets of OD results.

DRUG DURATION

Drug effects don't last forever. Eventually, the human body will filter out its blood supply, and normal functions will begin anew. The amount of time the drug lasts for determines the final base difficulty number. The difficulty number is used when a character attempts to create a drug from raw materials. Subsequently, it also helps determine the drug's base cost per dose, since drugs that are difficult to create are usually much more expensive than drugs you can produce using Mr. Wizard's Chemistry Kit.

DRUG DURATION	DIFFICULTY MULTIPLIER
Short - 1D10 Minutes	x1
Medium - 1D6×10 Minutes	x2
Long - 1D10 Hours	x3

To get the final difficulty number for a drug, add together the cost of its effects, its side effects, its strength, and multiply that sum by the difficulty modifier of the drug's duration. Voila! You're almost done! Remember that no drug's difficulty can be lower than 1, OR a negative value. If your final difficulty number is negative, drop the negative symbol and use the number as if it were a positive.

DRUG LEGALITY

Not all drugs are illegal. At the same time, though, not all drugs are available at Save-On drugs, either. So we come to the question of legality. Decide for yourself if the drug you're designing is something that the average vice squad would want to see on the street. Legality is important, because it helps determine the base cost of the drug in question.

LEGAL/COMMON

Legal/Common drugs include anything you can buy in a liquor store, bar, or over the counter in a pharmacy. It's common stuff that is easy to get, legal, but still governed by local and federal laws.

BY PRESCRIPTION ONLY

Anything available By Prescription Only includes substances that must be obtained through a doctor or pharmacist, prescribed for a specific ailment. If you've got a prescription, you can get the drug pretty easily, and for a much reduced cost. However, getting the stuff on the street may take a bit more doing, and cops don't like to find you carrying around a bottle of pills with someone else's name on it.

ILLEGAL (TYPE A, TYPE B, and TYPE C)

Illegal drugs are anything that the FDA hasn't approved of. Dangerous non-medicinal chemicals are also ruled by these categories, as are combat drugs. The local cops and feds try to stop these sorts of drugs from being sold on the streets, usually without much success. Illegal drugs are broken up into three categories: Type A, Type B, and Type C. Type C substances are illegal, but possession is usually a minor offense. Type B substances are usually more dangerous than Type C, and carry with them a stiffer punishment. Type A substances include the most dangerous and addictive drugs, and penalties for carrying them are large.

EXPERIMENTAL

Experimental drugs are corporate, criminal, and military cocktails that end up on the streets as part of clandestine experiments. Most of the time, they are part of a single shipment, and don't stay around too long unless they sell well. Drugs like this don't remain in this category for long, and are usually classified as an Illegal-rated substance after a few weeks on the streets.

BASE COST

The base cost in dollars (or standard monetary unit) of a drug is determined by multiplying the final difficulty number by the legality multiplier supplied on the chart below. The resulting number is the base cost for obtaining the substance on the street. The drug's final cost is determined by the form that it comes in.

DRUG LEGALITY	PRICE MULTIPLIER	EXAMPLE
Legal/Common	x1	Alcohol, Tobacco,
By Prescription Only	x5	Valium, Xanax, Sleeping Pills
Type C Illegal	x5	Marijuana,
Type B Illegal	x7.5	Cocaine, Crystal Meth,
Type A Illegal	x10	LSD, Heroin
Experimental	x25	?WHO KNOWS?



DRUG FORMS

The drug's form defines its physical characteristics and how the user will interface with it. Is it something you can smoke? Is it a pill? Do you use a syringe or an air hypo to inject it? The drug form also determines how much the final drug will cost, whether or not it can be "cut", and how fast it takes effect.

DRUG FORM	TIME TO EFFECT	COST MULTIPLIER	CAN IT BE CUT?
Pill, Tablet	2D6×10 Minutes	x 0.5	No
Gel Cap, Caplet	9+1D6 Minutes	x 1	No
Paper Tab	1D10 Minutes	x 1	No
Smoked, Inhaled	1D5 Turns	x 1	Yes
Powdered, Snorted	1D2 Minutes	x 1	Yes
Injected	1D5 Turns	x 1.5	Yes
Liquid	1D10 Minutes	x 1.5	Yes
Derm, Slap 10+2D10 Patch Seconds		x 2	No
Contact	1D5 Turns	x 2.5	No

DRUG FORM DESCRIPTIONS

PILL, TABLET

This is your typical pill. It comes in all shapes, sizes, and colors. Pills are generally chalky and taste bad unless coated. They are the cheapest form of drug administration since they take a horribly long time to go into effect.

GEL CAP, CAPLET

This includes drugs that are specially coated for ease of swallowing, and that are time-released for a quicker effect than tablets. Most people prefer this form to tablets.

PAPER TAB

A paper tab is a piece of paper that has been soaked in a concentrated liquid drug and then dried. The user places the tab onto his tongue and sucks on it until the drug takes effect. This form of drug has always been a popular way of administering hallucinogens like LSD.

SMOKED, INHALED

These drugs come in a form that is easily burnt. The smoke or fumes produced are then inhaled directly through a device, usually a pipe or electric vaporizer. Sometimes, tobacco or cigarettes are soaked in a liquid drug and then dried and smoked.

POWDERED, SNORTED

Powdered drugs that are snorted make their way to the lungs, sinuses, and mucus membranes where they are absorbed by the body. This form of use is less offensive to some than smoking, but it also takes a little longer for the drug to take effect.

INJECTED

Most injected drugs are concentrated liquids designed for direct entry into the bloodstream. Some start out as powdered drugs, but are liquefied in a process known as "free basing." Some people prefer this method of using, and will pay the extra cash to avoid the risks of freebasing the product themselves.

LIQUID

Liquid drugs are usually swallowed like a drink. Some can be mixed with normal soft drinks or fruit juices to cut the taste, though some are designed to be quite tasty all by themselves. While it may take the drugs longer to take effect, this is by far one of the most comfortable methods of imbibing.

DERM, SLAP PATCH

An adhesive sticker that is placed upon the skin, and the dose of the drug is then absorbed. Derms are convenient to use, easy to transport, and, if sealed in plastic, they have a good shelf life. However, the adhesive can irritate sensitive skin and cause an itchy rash.

CONTACT

Contact drugs are absorbed directly by the skin, almost instantly. They are usually hard to keep for long periods of time, as they are unstable and "go bad" quickly. However, they are the easiest and most comfortable way to administer drugs (or poisons) to yourself or an unsuspecting victim.

CUTTING DRUGS

are already cut with an inactive substance, since they would be far too powerful (and possibly deadly) if they weren't toned down. Of course, most dealers don't have your safety in mind when cutting their products. Unscrupulous dealers want to make their stock last a lot longer, so they cut it down and charge normal price. It is up to the individual user to check the quality of the dope he buys. A drug that is cut loses strength equal to the percentage it is

Drugs are "cut" for any number of reasons. Most powdered drugs

buys. A drug that is cut loses strength equal to the percentage it is cut by, rounding down. For example, a strength 4 euphoric that has been cut 25% with baby laxative will have a relative strength of 3, as opposed to 4 (its potency is reduced by 25% of the original STR). If cut 50%, it will only have a STR of 2, and so on.

FINAL COST DETERMINATION

Multiply the base Euro cost of the drug by the drug form cost multiplier (above) to find the drug's final cost per dose. This is an average price, based on the price of materials used and man-hours of work involved. Dealers will charge what the market will bear, however, and price increases during shortages and police crackdowns are all too common. On the up side, drugs which have a very low demand will cost significantly less. Final price is determined by the GM's discretion.

LABRATORIES

All Pharmaceuticals are made in labs, whether this lab is a corporate facility, or a mobile home kitchen, depends on the drug and the chemist. For game-play, Labs fall into four classifications. These classifications determine the base cost, difficulty levels of the drugs it can yield, and the number of batches it can produce at a time.

1: Simple - \$100 Diff 1-10 drugs.

This represents most home-made labs and stills found in basements and abandoned buildings. It is usually a jury-rigged assortment of parts, few of which being used for their intended purpose (soda bottle beakers, duct tape, etc). It requires a minimum Chemistry roll of 15 and 1 hour to assemble. If the Chemist rolls a 13 or 14, the lab will still work, but at +5 difficulties to Pharmaceutical checks. If the Chemist succeeds by 5 or more, the Pharmaceutical checks in the lab receive a +1 bonus.

Only drugs whose Difficulty is 10 or less can be manufactured in Simple labs. It is possible to create drugs up to 20 Difficulty, but with a +5 difficulty to Pharm rolls.

A Simple Lab can create only 1 drug batch at a time.

2: Moderate - \$1000 Diff 11-20 drugs.

This represents most high school-level chemistry labs. Equipment might not be the best, but it works better than home-made labs. It

requited a minimum Chemistry roll of 15 and 5 hours to assemble. If the chemist rolls a 13 or 14, the lab will still work, but at +5 difficulty to Pharm checks. If they succeed by 5 or more, Pharm checks receive a +1 bonus.

Only drugs with Difficulty of 20 or less can be manufactured in Moderate Labs. Drugs with difficulty 1-10 receive a +1 bonus to create. It is possible to create drugs up to 30 Difficulty, but with a +5 difficulty to Pharm rolls.

A Moderate Lab can create up to 2 batches of a single drug at a time.

3: **Advanced** - \$5,000 Diff 21-30 drugs

This represents most College or small professional laboratories. These labs are non-mobile and often require modifications to the structure they are housed in. It requires a minimum Chemistry roll of 20 and 20 hours to assemble. If the Chemist misses their roll by less than 5, it will still work, but at a +5 difficulty to all Pharm checks. If the Chemist succeeds by 5 or more, Pharm checks receive a +1 bonus.

Only drugs with Difficulty 30 or less can be manufactured in Advanced labs. Drugs with Difficulty below 10 receive a +3 bonus to create. Drugs with Difficulty 11-20 receive a +1 bonus to create. It is possible to create drugs whose difficulty is above 30, but at a +5 difficulty to Pharm Rolls

An Advanced Lab can create up to 5 batches of a single drug at a time.

4: Sophisticated - \$25,000 Diff 31+ drugs

This represents only the most sophisticated of laboratories. Much of its equipment is unique and the facility it's housed in is built to the lab's specific needs. Often including independent air with nano-filtration systems, and robotic manipulators. When taking into consideration the costs of the tailored facility housing the lab, this can cost in the hundreds of millions. It requires a minimum Chemistry roll of 25 and 180 hours to assemble. If the Chemist misses their roll by less than 5, it will still work, but at a +5 difficulty to Pharm checks. If the Chemist succeeds by 5 or more, Pharm checks receive a +1 bonus

Any drug can be created in Advanced labs. Drugs with Difficulty below 10 receive a +5 bonus to create. Drugs with Difficulty 11-20 receive a +3 bonus to create. Drugs with Difficulty 21-30 receive a +1 bonus to create.

An Advanced Lab can create up to 8 batches of a single drug at a time.

Manufacturing Costs:

drugs.

Creating a drug requires supplies that are used every time the creation is attempted. These costs fall into 2 categories: Fixed Cost, and Batch Cost.

Fixed Cost: This represents the cost of tailoring a lab for the specific drug being produced. Only one drug can be produced at a time, and this cost must be paid every time the lab is configured for a specific drug. The fixed cost is the Drug's Difficulty x20. This cost need only be paid once per drug type per lab. Configuring a Lab to a specific drug will require a Chemistry roll greater than the difficulty of the drug to be made (minimum 10). This roll is made every time the lab is configured, even if the monetary cost no longer has to be paid for previously-created

Batch Cost: This represents the base cost 1 batch of a specific drug costs. This includes chemicals, and materials consumed in the manufacturing process. The batch cost is the Drug's Difficulty x5, plus any Cost Multipliers (see DRUG FORMS). This money is

spent every attempt to produce a drug, whether the Pharmacist was successful or not. Example: Killroy is a DC16 Drug that is injected. Every attempted batch will cost (16x5x1.5) \$120 to create. That's a lot, but considering each dose of Killroy is \$180, the cost is made up very quickly.

Manufacturing Time and Volume:

Time: It takes 1 hour for every point of difficulty to create a single batch of a drug. For example, Killroy, which has a Difficulty 16 to create, will take a minimum of 16 hours to manufacture 1 batch.

Volume: A single batch can produce 2D6 + the Pharmacist's level in both Pharmacy Tech and Chemistry, in doses. For Example: a character with a Med: Pharmaceutical Tech of 5, Chemistry 6 can make 2D6+11 doses with every batch.



DRUG LAB 101 QUICK REFERENCE CHARTS

DRUG EFFECT	Cost	DRUG EFFECT	Cost
Antibiotic	10	Anticoagulant	10
Antidote	15	Aphrodisiac	10
Attribute Increase	20	Coagulant	10
Contraceptive	10	Depressant	5
		•	5
Enhanced Perception	15	Euphoric	
Hallucinogen	10	Hypnotic	15
Increased Endurance	10	Increased Healing Rate	15
Pain Negation	10	Soporific	5
Stun Reducer	10	Toxin: Bio	10
Toxin: Neuro	15		
DRUG FEATURE	Modifier	DRUG FEATURE	Modifier
Lessened Withdrawal Symptoms	+10	Longer Duration	+10
Timed Duration	+10		
UNTIMED SIDE EFFECT	Modifier	UNTIMED SIDE EFFECT	Modifier
Addiction Tolerance	-4	Carcinogenic	-10
Cardiac Arrest	-12	Coma	-10
Death	-15	Genetic Damage	-6
Nerve Degeneration	-15	Nightmares	-6
Permanent Sense Loss	-6	Physiologically Addictive	-5/-10/-15
Possible Attribute Loss	-8	Psychologically Addictive	-4/-8/-12
Severe Withdrawal Symptoms	-6	Sterility	-8
TIMED SIDE EFFECT	Modifier	TIMED SIDE EFFECT	Modifier
Acne	-2	Aggressive Behavior	-6
Bad Breath	-2	Bloodshot Eyes	-2
Cold Sweats	-2	Constipation	-2
Cowardice	-6	Dandruff/Eczema	-2
Dehydration	-2	Delusions	-5
Depression	-4	Diarrhea	-4
Difficult Respiration	_	D: .:	
Diggings	-6	Diuretic	-2
Dizziness	-6 -5	Easily Bruised	-2 -2
Excessive Salivation			
	-5	Easily Bruised	-2
Excessive Salivation	-5 -2	Easily Bruised Flatulence	-2 -2
Excessive Salivation Hair Loss	-5 -2 -2	Easily Bruised Flatulence Hallucinations	-2 -2 -5
Excessive Salivation Hair Loss Headaches	-5 -2 -2 -4	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion	-2 -2 -5 -2
Excessive Salivation Hair Loss Headaches Hives	-5 -2 -2 -4 -4	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity	-2 -2 -5 -2 -2
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding	-5 -2 -2 -4 -4	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear	-2 -2 -5 -2 -2 -10
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy	-5 -2 -2 -4 -4 -4 -3	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure	-2 -2 -5 -2 -2 -10 -8
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration	-5 -2 -2 -4 -4 -4 -3 -5	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity	-2 -2 -5 -2 -2 -10 -8 -2
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite	-5 -2 -2 -4 -4 -4 -3 -5 -2	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition	-2 -2 -5 -2 -10 -8 -2 -4
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss	-5 -2 -2 -4 -4 -4 -3 -5 -2 -4	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea	-2 -2 -5 -2 -10 -8 -2 -4 -4
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss Numbness	-5 -2 -4 -4 -4 -3 -5 -2 -4 -4 -4 -4 -5 -5 -2 -4 -4/-6	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea Paralysis	-2 -2 -5 -2 -10 -8 -2 -4 -4 -6
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss Numbness Paranoia	-5 -2 -4 -4 -4 -3 -5 -2 -4 -4/-6 -5	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea Paralysis Psychotic Rage	-2 -2 -5 -2 -10 -8 -2 -4 -6 -10
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss Numbness Paranoia Reduced Attribute	-5 -2 -4 -4 -4 -3 -5 -2 -4 -4/-6 -5 -5	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea Paralysis Psychotic Rage Reduced Awareness	-2 -2 -5 -2 -10 -8 -2 -4 -4 -6 -10 -6
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss Numbness Paranoia Reduced Attribute Ringing In Ears	-5 -2 -4 -4 -4 -3 -5 -2 -4 -4/-6 -5 -5 -2	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea Paralysis Psychotic Rage Reduced Awareness Runny Nose	-2 -2 -5 -2 -10 -8 -2 -4 -4 -6 -10 -6 -2
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss Numbness Paranoia Reduced Attribute Ringing In Ears Sense Reduction	-5 -2 -4 -4 -4 -3 -5 -2 -4 -4/-6 -5 -5 -2 -4	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea Paralysis Psychotic Rage Reduced Awareness Runny Nose Sexual Arousal	-2 -2 -5 -2 -10 -8 -2 -4 -6 -10 -6 -2 -4
Excessive Salivation Hair Loss Headaches Hives Internal Bleeding Itchy Lack of Concentration Loss of Appetite Memory Loss Numbness Paranoia Reduced Attribute Ringing In Ears Sense Reduction Sleepy	-5 -2 -4 -4 -4 -3 -5 -2 -4 -4/-6 -5 -5 -2 -4 -4	Easily Bruised Flatulence Hallucinations Heartburn/Acid Indigestion Impotence/Frigidity Irrational Fear Kidney/Liver Failure Light Sensitivity Loss of Inhibition Nausea Paralysis Psychotic Rage Reduced Awareness Runny Nose Sexual Arousal Strange Skin Pallor	-2 -2 -5 -2 -10 -8 -2 -4 -4 -6 -10 -6 -2 -4 -2

Note: All timed side-effects can be bought as "Possible Side Effects" at half cost.

DRUG STRENGTH	COST	DURATION	MULTIPLIER
Strength +1	1	Short - 1D10 Minutes	x1
Strength +2	2	Medium - 1D6×10 Minutes	x2
Strength +3	4	Long - 1D10 Hours	х3
Strength +4	8		
Strength +5	16		

DRUG LEGALITY	PRICE MULTIPLIER
Legal/Common	x1 Dollars
By Prescription Only	x5 Dollars
Type C Illegal	x5 Dollars
Type B Illegal	x7.5 Dollars
Type A Illegal	x10 Dollars
Experimental	x25 Dollars

DRUG FORM	TIME TO EFFECT	COST MULTIPLIER	CAN IT BE CUT?
Pill, Tablet	2D6×10 Minutes	x 0.5	No
Gel Cap, Caplet	9+1D6 Minutes	x 1	No
Paper Tab	1D10 Minutes	x 1	No
Smoked, Inhaled	1D5 Turns	x 1	Yes
Powdered, Snorted	1D2 Minutes	x 1	Yes
Injected	1D5 Turns	x 1.5	Yes
Liquid	1D10 Minutes	x 1.5	Yes
Derm, Slap Patch	10+2D10 Seconds	x 2	No
Contact	1D5 Turns	x 2.5	No

EQUATIONS

$$\label{eq:total_problem} \begin{split} Total \ Difficulty = Effects + Side \ Effects + Drug \ STR \ x \ Duration \\ Multiplier \end{split}$$

Base Cost = Total Difficulty x Legality Multiplier (see VARIABLE DRUG COSTS)

Final Price = Base Cost x Price Multiplier (see DRUG FORMS)

MANUFACTURING:

Fixed Cost = Total Difficulty x 20
Batch Cost = Total Difficulty x 5 x Price Multiplier
(see DRUG FORMS)

of Doses Per Batch = Chemistry Level + Pharm Level + 2D6 Time to Produce = (Total Difficulty) hours

Lab Type	Cost	Build Difficulty	Hours to Assemble	Batches at a Time
Basic	\$100	15	1	1
Moderate	\$1,000	15	5	2
Advanced	\$5,000	20	20	5
Sophisticated	\$25,000	25	180	8

Lab Type	Drug Level 1-10	Drug Level 11-20	Drug Level 21-30	Drug Level 31+
Basic	+0	-5	n/a	n/a
Moderate	+1	+0	-5	n/a
Advanced	+3	+1	+0	-5
Sophisticated	+5	+3	+1	+0

(Created and written for Cyberpunk 2020 by Gary Ostlelford (ocelot@cox.net),
Additional materiel written by Seth Skorkowsky
Edited for Interlock Unlimited by Deric Bernier (droc@mc2k.com)
All images by Deric Bernier.

For more of the best Interlock Unlimited goodness available, go to



